



UW SOIL AND FORAGE
LAB

YOUR UW

SOIL PLANT TISSUE FORAGE MANURE LIME RESEARCHERS CONTACT
FEES/BROCHURE PAYMENT

Contact & location:

4702 University Avenue
Madison, WI 53705
(608) 262-4364
soil-lab@mailplus.wisc.edu

Soil > Wildlife Food Plots

WILDLIFE FOOD PLOTS

Routine Analysis: \$15 per sample. Includes pH, lime requirement, organic matter, phosphorus, potassium, and nutrient recommendations.

Optional: Calcium + Magnesium: \$3 | Boron: \$3 | Zinc: \$3 |
Manganese: \$3 | Sulfur-Sulfate: \$3

Forms & Instructions:

Submission form – please use crop code numbers from Form
Instructions / Crop Codes

Instructions for Sampling Field Soils (A2100)

Soil Names (Excel) | A2809 Soil map units and yield potentials
(12/15/2015)

How to Determine Soil Map Units

Analysis and Reporting

Your soil test report will provide an accurate index of the level of available nutrients in the soil; indicate the degree of nutrient deficiency that may exist for the various crops grown; suggest how the deficiency might be corrected; and provide the results in a meaningful way so that you can make the appropriate decision as to what nutrients to add. View a sample report.

Laboratory analysis will include soil pH, percent organic matter, phosphorus and potassium. Additional secondary and micro-nutrient analysis of calcium, magnesium, boron, manganese, zinc and sulfur are also available for an additional charge.

Additional References & Resources

- Nutrient Application Rate Guidelines for Field, Vegetable, and Fruit Crops in Wisconsin (A2809)
- Sampling Soils for Testing (A2100)
- Management of Wisconsin Soils (A3588)
- Nitrogen Application Rate Guidelines for Corn (MRTN)
- Supplemental Nitrogen to Corn Price Ratio Calculator
- Wisconsin Soil Test Summary Data

- SnapPlus

©2022 Board of Regents of the University of
Wisconsin System

Feedback, questions or accessibility issues:

websupport@cals.wisc.edu

[Log In](#)

[Request Help](#)

[Help Docs](#)



Wildlife Food Plot Soil Sample Information Form

For Lab Use Only: Date: Lab No:	Please check how you would like to receive your results: <input type="checkbox"/> U.S. Mail <input type="checkbox"/> Email	Method of Payment:
	Name:	Account ID
	Address:	OR Amt. Paid \$
	City: State: Zip:	<input type="checkbox"/> Cash
	Phone:	<input type="checkbox"/> Check No.
	Email:	<input type="checkbox"/> Card (we will call)

TOTAL NUMBER OF SAMPLES: _____	PLOW DEPTH: _____	Complete form instructions can be found on the back of this sheet. Use a separate submission form for samples from different counties.
---------------------------------------	--------------------------	---

SAMPLE #	SAMPLE NAME	COUNTY <small>(where samples were taken)</small>	SOIL NAME <small>(REQUIRED)</small>	Acres in Field	Slope %	4-YEAR CROP ROTATION	Previous Legume Crop codes below) <small>(see</small>
						Sequence to be Grown <small>(crop code)</small>	Legume Crop <small>(crop code)</small>
1							
2							
3							
4							
5							
6							

Wildlife Food Plot (4-YR Rotation) Crop Codes:	Previous Legume Crop Codes (for nitrogen credit):
69 Corn/brassicas 70 Legume-grass mix 71 Oat/wheat/rye 72 Soybean 73 Sugar beet/turnip	1 Alfalfa or red clover established one year or less 2 Alfalfa or red clover established one year or more 3 Legume-grass pasture or hay field 4 Soybeans 5 Peas, beans or other leguminous crop

Tests include: pH, lime requirement, organic matter, phosphorus (P) and potassium (K). <div style="text-align: center;">Special Soil Tests (for an additional fee) <small>(List field or sample number)</small></div> <table style="width: 100%; border-collapse: collapse;"> <tr><td style="width: 50%;">Calcium/Magnesium</td><td style="width: 50%;">Zinc</td></tr> <tr><td>Boron</td><td>Sulfate</td></tr> <tr><td>Manganese</td><td>Other</td></tr> </table>	Calcium/Magnesium	Zinc	Boron	Sulfate	Manganese	Other	SOIL NAME Write the full name (NOT the abbreviation) from an FSA farm plan or NRCS soil survey map. 4-YEAR CROP ROTATION Indicate the intended crops to be grown for the next four years (one crop code per year, from the WFP (4-YR Rotations Crop Code box only). <div style="text-align: right;">\$15.00/sample</div>
Calcium/Magnesium	Zinc						
Boron	Sulfate						
Manganese	Other						
Soil tests recommended if: growing legume forage B and SO₄-S growing corn Zn and SO₄-S growing small grain or soybean (with soil pH >7.0) Mn acid or sandy soil with high amounts of applied K Ca/Mg	1/22						

INSTRUCTIONS

CHECK how you wish to receive your results. **PRINT YOUR NAME - Please Print clearly**

Fill in the county from which the sample(s) were taken. Please fill out separate forms for **each** county. Fill in your address and the e-mail address if you would like the results emailed to you.

METHOD OF PAYMENT

Fill in your Account ID **or** your payment (cash, credit card, or check payable to: **UW Soil Testing Lab**).

Wildlife Food Plot is \$15.00/sample.

SOIL NAME

Write the **full** soil name (**NOT** the abbreviation) from an FSA farm plan or NRCS survey map (can be found online).

Example: Fayette silt loam, write "Fayette". If the field has more than one soil type, use the most predominant soil found in the field. A more precise soil test recommendation can be given if the soil name is included. If soil name is unknown, please be sure to include the county where the soil samples were taken.

4-YEAR CROP ROTATION

Indicate the intended crops to be grown for the next four years (one crop code per year). Use crop codes from the Wildlife Food Plot (4-YR Rotation) Crop Codes box only.

PREVIOUS LEGUME CROP

Enter the crop code for the previous legume crop grown on the field (choose codes from the "Previous Legume Crop Codes" box only). One code per sample.

SPECIAL SOIL TESTS: Special tests may be run on individual samples. Please list samples numbers next to special tests to be run. Special soil tests are \$3.00 per test, per sample.

INSTRUCTIONS

Madison Lab: 608-262-4364

Marshfield Lab: 715-387-2523

CHECK how you wish to receive your results. **PRINT YOUR NAME - Please Print clearly**Fill in the county from which the sample(s) were taken. Please fill out separate forms for **each** county. Fill in your address and the e-mail address if you would like the results emailed to you.**METHOD OF PAYMENT**Fill in your Account ID **or** your payment (cash, credit card, or check payable to: **UW Soil Testing Lab**)**FIELD ID and SAMPLE NUMBER(S)****Please number samples consecutively.**

Record the field and sample identification for each field on the same line.

EXAMPLE

Field ID	Sample No(s)
1	1-4
2	5
3	6-8

SOIL NAME [REQUIRED]Write the **full** soil name (**NOT** the abbreviation) from an FSA farm plan or county soil survey map.

Example: Fayette silt loam, write "Fayette". If the field has more than one soil type, use the most predominant soil found in the field. A more precise soil test recommendation can be given if the soil name is included.

4-YEAR CROP ROTATIONIndicate the intended crops to be grown for the next four years (**one** crop code per year).**Use the crop codes(s) listed in the table below.**

Enter a yield goal no more than 10-15% higher than the prior 5-year average for each crop. Base the yield goal for corn on yield of No. 2 corn at 15.5% moisture. Yield goal for alfalfa should be based on dry matter in T/a. Base yield of other crops on the yield unit shown in parenthesis (). Give yield goals to the nearest ½ T for crop units expressed in T/a.

Crop Code	Crop Name	Yield Unit	Crop Code	Crop Name	Yield Unit	Crop Code	Crop Name	Yield Unit
1	Alfalfa, established	(ton)	66	CRP, alfalfa	(n/a)	40	Pumpkin	(ton)
2	Alfalfa, seeding	(ton)	68	CRP, grass	(n/a)	64	Raspberry, estab.	(all)
60	Apple, establishment	(all)	67	CRP, red clover	(n/a)	76	Rye, grain	(bu)
3	Asparagus	(lb)	20	Cucumber	(ton)	43	Rye, grain + straw	(bu)
74	Barley, grain	(bu)	21	Flax	(bu)	82	Sm grain & legume silage	(ton)
4	Barley, grain + straw	(bu)	22	Ginseng	(lb)	30	Sm grain/leg slg, w/alf sdg	(ton)
5	Bean, dry (kidney, navy)	(cwt)	79	Grape, establishment	(all)	81	Small grain silage	(ton)
6	Bean, lima	(lb)	84	Grass, hay	(ton)	29	Sm grain silage, w/alf sdg	(ton)
44	Bean, snap	(ton)	41	Grass, reed canarygrass	(ton)	46	Sorghum, grain	(bu)
7	Beet, table	(ton)	45	Grass, sod for turf, est.	(all)	47	Sorghum-sudan, forage	(ton)
61	Blueberry, estab.	(all)	85	Grass, switchgrass	(ton)	48	Soybean, grain	(bu)
8	Brassica, forage	(ton)	86	Hop	(lb)	77	Soybean, grain + straw	(bu)
9	Broccoli	(ton)	23	Lettuce	(ton)	49	Spinach	(ton)
10	Brussels sprouts	(ton)	24	Lupine	(bu)	50	Squash	(ton)
11	Buckwheat	(lb)	25	Melon	(ton)	65	Strawberry, estab.	(all)
12	Cabbage	(ton)	26	Millet	(bu)	51	Sunflower	(lb)
13	Canola	(bu)	27	Mint, oil	(lb)	52	Tobacco	(lb)
14	Carrot	(ton)	75	Oats, grain	(bu)	53	Tomato	(ton)
15	Cauliflower	(ton)	28	Oats, grain + straw	(bu)	54	Trefoil, birdsfoot	(ton)
16	Celery	(ton)	31	Onion	(cwt)	55	Triticale, grain	(lb)
62	Cherry, establishment	(all)	34	Past.,legume(<30%)-grass	(ton)	80	Triticale, grain + straw	(lb)
42	Clover, red	(ton)	83	Past.,legume(>30%)-grass	(ton)	56	Truck crops	(all)
17	Corn, grain	(bu)	33	Pasture, grass	(ton)	57	Vetch, crown/hairy	(ton)
38	Corn, popcorn	(bu)	32	Pasture, unimproved	(ton)	78	Wheat, grain	(bu)
18	Corn, silage	(ton)	35	Pea, canning	(lb)	58	Wheat, grain + straw	(bu)
19	Corn, sweet	(ton)	36	Pea, chick/field/cow	(ton)			
63	Cranberry, estab.	(all)	37	Pepper	(ton)			
			39	Potato	(cwt)			

FERTILIZER CREDIT INFORMATION: Legume-sod plowdown or manure application may reduce nutrient need.**Previous Legume Crop:** Enter the crop code for the previous legume crop grown on the field. For all forage crops that were plowed down, indicate the % legume remaining in stand and check if there was more than 8 inches of regrowth in the fall before the stand is killed.**Manure Applied to Field Since Last Crop:** If manure was applied to the field since harvesting the last crop, choose manure code from **Manure Code List** on front of Information Sheet. Specify the approximate rate of application in T/a for solid or 1000 gal/a for liquid manure, time to incorporation and the number of consecutive years manure has been applied to this field.**SPECIAL SOIL TESTS:** Special tests may be run on individual samples, or all the samples from the same field may be combined at the lab for a single field analysis. If the special test(s) is requested on a field basis only, enter the field ID.If the special test(s) is requested for each sample, enter the field ID and sample number.

Sampling soils for testing

John B. Peters and Carrie A.M. Laboski

A soil test is the only practical way of determining whether lime and fertilizer are needed for a specific crop. However, if a soil sample does not represent the general soil conditions of the field, the recommendations based on the sample may be misleading. An acre of soil to a 6-inch depth weighs about 1,000 tons, yet less than 1 ounce of soil is used for each test in the laboratory. Therefore, it is very important that the soil sample be representative of the entire field.

Before collecting soil samples, you should determine the overall approach of the nutrient management program. This will affect the number of samples needed and method by which samples will be taken. Specifically, will nutrient and lime applications be made at a single uniform rate for the whole field being tested or will applications be made at variable rates to field areas that have been identified as having different soil test levels?

Goals of a soil sampling program

When sampling soils for testing and obtaining fertilizer and lime recommendations, the most common objectives are to:

1. Obtain samples that accurately represent the field from which they were taken.
2. Estimate the amount of nutrients that should be applied to provide the greatest economic return to the grower.
3. Estimate the variation that exists within the field and how the nutrients are distributed spatially.
4. Monitor the changes in nutrient status of the field over time.

Selecting a soil sampling strategy

Before selecting a sampling strategy, consider analytical costs, time and equipment available, field fertilization history, and the likelihood of a response to applied nutrients.

Sampling fields for a single whole field (uniform) recommendation

With conventional sampling, you will receive a single set of nutrient and lime application guidelines that are based on sample averages. The sampling guidelines in Table 1 are based on when a field was last tested (more or less than 4


Table 1. Recommended sample intensity for uniform fields.

Field characteristics	Field size (acres)	Suggested number of samples ^a
Fields tested more than 4 years ago OR fields testing in the responsive range	All fields	1 sample/5 acres
Nonresponsive fields tested within past 4 years	5–10	2
	11–25	3
	26–40	4
	41–60	5
	61–80	6
	81–100	7

^a Collect a minimum of 10 cores per sample.


years ago) and whether the field was responsive or nonresponsive the last time it was tested. The field is considered to be in the responsive range if either soil test phosphorus (P) or potassium (K) levels are in the high (H) category or lower. A nonresponsive field is one where both soil test P and K levels are in the very high (VH) or excessively high (EH) categories.

Each sample should be made up of a minimum of 10 cores to ensure accurate representation of the nutrient needs of the field. Research has shown that taking 10 to 20 cores provides a more representative sample of the area than when samples are made up of fewer cores. When gathering soil cores to make a composite sample, use a W-shaped sampling pattern (as shown in Figure 1) over the whole area the sample represents. Be sure to thoroughly mix the cores before placing approximately 2 cups in the sample bag.



For best results, submit multiple samples for all fields. When at least three samples are provided for a field, samples that are significantly higher than the field average may be discarded and an adjusted average calculated. Using an adjusted

Figure 1. Recommended W-shaped sampling pattern for a 15-acre field. Each sample should be composed of at least 10 cores.



average helps ensure that no part of the field is under-fertilized.

Where only one or two samples are taken in a field, no sample will be discarded, whereas one sample can be discarded if three or four samples are taken, and up to two samples may be discarded from fields having five or more samples. The criteria that determine if soil samples should be omitted from the field average include:

- If the average soil test P for a field is 35 parts per million (ppm) or less, samples that exceed the field average by more than 5 ppm may be removed and the field average recalculated.
- If the field average is greater than 35 ppm P, no samples will be discarded.
- If the average soil test K for a field is 175 ppm or less, samples that exceed the field average by more than 20 ppm may be discarded and the field average recalculated.
- If the field average is greater than 175 ppm K, no samples will be discarded.

It is not appropriate to vary nutrient application rates across sampling areas when using the whole field (uniform) soil sampling scheme.

Sampling fields for site-specific management


Site-specific management requires a distinct picture of the magnitude and location of soil test variability. Sampling soils for site-specific management usually involves taking many more composite samples than sampling for a single recommendation. A global positioning system (GPS) is used to record the geographical coordinates of each sample. This information is used to generate an application map by using various mathematical techniques to interpolate the nutrient application rate between sampling points. Using variable rate application technology, these fields can be managed more intensively than the conventional approach of one fertilizer and lime rate per field. A careful evaluation of the economics of this intensive

of a sampling system needs to be done before proceeding.

When using a site-specific approach to soil sampling, sample handling and testing are similar to the traditional system, but recommendations may vary from one part of the field to another, and these areas must be managed separately to realize the potential advantages of intensive soil sampling.

Several sampling strategies can be used to guide variable-rate fertilizer and lime applications. Grid sampling uses a systematic approach that divides the field into squares of approximately equal size (grid cells). The sampling technique used is known as **grid-point sampling**. A grid-point sample consists of at least 10 cores collected from a small area (10-foot radius) around a geo-referenced point. When using a grid sampling approach, Wisconsin research recommends a sampling strategy based on an unaligned systematic grid (Figure 2). Sampling points should be unaligned because sampling in a uniform grid arrangement may lead to biased results if aligned with row patterns. Fields that have soil test P and K levels in the nonresponsive categories should be grid-point sampled on a 300-foot grid. This is equivalent to one soil sample for every 2 to 2.5 acres. Where there is no information about the P or K status of the field or where previous tests were in the responsive range, a 200-foot grid size should be used. This is equivalent to approximately one soil sample per acre. Wisconsin research indicates these small grid cell sizes are needed to adequately characterize the variability in soil fertility. A larger grid cell size (such as 5 acres) may not adequately describe the field variability and may

Figure 2. An example of an unaligned grid pattern for grid sampling fields.



limit the potential economic benefits of site-specific management.

Other considerations in selecting a sampling strategy

Select the sampling strategy appropriate for the field size and topography.

Contour strips. On contour strip fields, sample each strip separately if it is approximately 5 acres or more in size, following the sampling intensity guidelines provided in Table 1. Cores from two or three small strips that have identical cropping and management histories may be combined following these same recommended sampling intensity guidelines. Using a grid-point sampling approach on contour strips or small fields is not appropriate, regardless of grid cell size. This is because a grid technique may result in many soil samples being collected from one contour strip but none in other strips; additionally, grid-point samples may be on the edge of the strips and not adequately represent the strip.

Five-acre grid-point sampling. The 5-acre grid point sampling system for whole field management recommendations has recently become popular with soil samplers because it takes less time to collect cores, compared to the traditional W pattern. Another advantage of this approach is its ability to track changes in soil test levels over time, because soil samples are collected from the same geo-referenced point each time the field is sampled. Five-acre grid-point sampling can likely be used in some situations and not in others. For example, in fields that were soil sampled within the past 4 years and tested in the nonresponsive range, averaging the soil test results from 5-acre grid-point sampling is reasonable. This is because there previously had not been a fertilizer recommendation on these fields and some variability at excessively high soil test levels does not change the fact that no fertilizer was recommended. For fields that were sampled more than 4 years ago or where past soil test results were in the responsive range, 5-acre grid-point sampling may not be the best choice of sampling techniques. This is because 5-acre grid-point sampling may not adequately represent the variability within a field, and a comparatively small change in soil test level of 5 to 10 ppm could mean a large change in the amount of nutrients recommended. For

small fields and contour strips, taking a few 5-acre grid-point samples in each field and averaging them likely does not provide a representative sample of the field. Additionally, the total number of samples may be so few that none of them can be eliminated from the field average if it appears one is an outlier.

Smart (zone or directed) sampling.

Another approach gaining support among researchers is smart sampling, also known as directed or management zone sampling. This approach uses information that has been collected using other precision agricultural technologies such as yield maps, aerial photographs of bare soil or crop canopy, or soil electrical conductivity measurements. Directed sampling evaluates the spatial distribution of several factors that may influence nutrient availability and crop productivity to help define sampling areas with similar characteristics. With previous comments in mind, either the W pattern or grid-point method can be used to collect samples within management zones. If the results of grid or management zone sampling do not warrant variable-rate application (for example, relatively little between-sample variation), average them to determine the appropriate single-rate treatment.

Procedures for taking soil samples

When to take soil samples

Take soil samples at any convenient time. Studies examining the effect of sampling time on soil test results suggest that test values for pH and phosphorus (P) are typically slightly higher in early spring samples than in fall samples. The effect of time of sampling on soil test potassium (K) results is dependent upon clay mineralogy and soil test level. Soil test K results may be higher in spring compared to fall on lower testing soils, but on higher testing soils, soil test K may be lower in spring compared to fall. To receive your recommendations early enough to enable you to apply the lime and fertilizer needed, it may be best to sample in the fall. Another benefit of fall testing is that fertilizer prices are more likely to be discounted then. Hayfields can be sampled after any cutting. Regardless of when you sample, it is best to be consistent from one year to the next.

Winter sampling, or sampling when the soil is frozen, is permissible only when it is possible to take a uniform boring or core of soil to the appropriate depth. This may require using a portable power boring tool. Using a pick or spade to remove a few chunks of frozen soil from the surface will give inaccurate results.

How to take soil samples

Certain government agency programs require nutrient management plans prepared according to the current USDA-NRCS nutrient management standard (590). Soil sampling and testing procedures and nutrient application rates based on these soil tests must be consistent with the provisions of the 590 standard to be eligible for many cost-sharing programs. These provisions currently include: following the soil sampling techniques outlined above, soil testing by a Wisconsin certified laboratory, and use of nutrient application rates consistent with the guidelines contained in the University of Wisconsin-Extension publication *Nutrient Application Guidelines for Field, Vegetable, and Fruit Crops in Wisconsin* (A2809).

When ready to sample, use a sampling probe or auger. You can obtain these tools on loan from most county Extension offices (counties.uwex.edu) or fertilizer dealers. Avoid sampling the following areas:

- Dead furrows or back furrows
- Lime, sludge, or manure piles
- Animal droppings
- Near fences or roads
- Rows where fertilizer has been banded
- Eroded knolls
- Low spots
- Where stalks or large bales were stacked
- Headlands

In addition, avoid sampling areas that vary widely from the rest of the field in color, fertility, slope, texture (sandy, clayey, etc.), drainage, or productivity. If the distinctive area is large enough to receive lime or fertilizer treatments different from the rest of the field, sample it separately.

These steps will help you take full advantage of the Wisconsin nutrient applica-

tion guidelines and must be followed to be consistent with the 590 standard.

1. If manure or crop residues are on the surface, push them aside to keep from including them in the soil sample.
2. Insert the probe or auger into the soil to plow depth or at least 6 inches. The sampling depth should be consistent. To aid year-to-year comparisons, it is important to take repeated samplings from the same field to exactly the same depth.
3. Take at least 10 soil cores or borings for each composite sample and, preferably, at least two composite samples for every field. For nonresponsive fields greater than 5 acres in size, obtain, at a minimum, the number of samples specified in Table 1. For responsive fields, as well as all fields that have not been sampled in the past 4 years, take one composite sample for every 5 acres.
4. Thoroughly mix the sample, then place about 2 cups of soil in a sample bag.
5. Identify the bag with your name, field identification, and sample number.
6. Record the field and sample location on an aerial photo or sketch of the farm and retain for your reference. Record the GPS coordinates, if available.
7. Fill out the soil information sheet. A completely and carefully filled out information sheet will provide the most accurate nutrient recommendations.

Always include a soil test information sheet when submitting soil samples to a laboratory for testing. The soil test information sheet used by the UW Soil Testing Laboratories can be found at: http://uwlab.soils.wisc.edu/files/forms/rfs_front.pdf.

Provide the soil name and field history whenever possible for more accurate recommendations. Information about legume crops previously grown on the soil and manure application history is essential for proper nutrient crediting from these sources. Include soil names and/ or map unit symbols from county soil

survey reports, web soil survey (<http://websoilsurvey.nrcs.usda.gov/app/>), or individual farm conservation plans. To obtain this information, contact your county Extension agent, NRCS district conservationist, or the County Land Conservation Department (LCD).

How often to sample

Most fields should be retested at least every 4 years to monitor soil fertility levels of immobile nutrients and pH to prevent nutrient deficiencies and avoid excess nutrient accumulation. Crop nutrient removals over a 4-year period in most cropping systems will not change soil test levels enough to affect recommended nutrient application rates. Exceptions include sands and loamy sands, which should be tested every 2 years. Also, depending on the initial soil test P and K levels, cropping systems such as high-yielding corn silage or alfalfa may require more frequent testing to adequately monitor changes in soil test levels.

What to do with soil samples

The soil samples and a completed soil information sheet can be taken to your county Extension office for forwarding to a certified soil testing laboratory, sent directly to the soil testing laboratory, or delivered in person.

To receive nutrient application rate guidelines consistent with those found in A2809, submit your soil samples to one of the Wisconsin certified laboratories. The College of Agricultural and Life Sciences, University of Wisconsin–Madison and the University of Wisconsin–Extension, through the Department of Soil Science, operate soil testing laboratories at Madison and Marshfield. Several private laboratories are also certified, and are listed at <http://uwlab.soils.wisc.edu/wdatcp/>. To become certified, laboratories must use the soil testing methods and nutrient application rate guidelines specified by WDATCP and must also meet quality control standards through periodic analysis of quality control soil samples.

To have your soil tested by the University of Wisconsin, send your samples to either of the listed laboratories. Find a sample submission form at <https://uwlab.soils.wisc.edu/farm-soil/>.

Tillage system considerations when sampling

Moldboard plowing. Sample to the depth of tillage.

Chisel plowing and offset disking. Take soil samples to $\frac{3}{4}$ of the tillage depth. When possible, take soil samples before spring or fall tillage. Sampling before tillage lets you determine the sampling depth more accurately and avoid fertilizer bands applied for the previous crop.

Till-plant and ridge tillage. Sample ridges to a 6-inch depth and furrows (between rows) to a depth of 4 inches. Combine equal numbers of soil cores from ridges and furrows to make up the composite sample.

No-till. Fields that have not been tilled for 5 or more years may develop an acid layer on the surface from the use of nitrogen fertilizer. This acid layer could reduce the effectiveness of triazine herbicides. Unincorporated phosphorus (P) and potassium (K) are also likely to build up in the surface soil. If an acid layer is suspected, take a separate sample to a depth of only 2 inches. When sending the soil to the lab, indicate that the sampling depth was only 2 inches. This sample will be tested for pH only, unless P and K are specifically requested. For fertilizer recommendations, take a separate sample to a depth of 6 to 7 inches. Fertilizer recommendations require this sampling depth because fertilizer calibration studies are based on plow-depth sampling. Sample between rows to avoid fertilizer bands.



Soil and Plant Analysis Laboratory

8452 Mineral Point Road
Verona, WI 53593-8696
(608) 262-4364

Soil and Forage Analysis Laboratory

2611 Yellowstone Drive
Marshfield, WI 54449-8401
(715) 387-2523

UW
Extension
Cooperative Extension

© 2013 University of Wisconsin System Board of Regents and University of Wisconsin–Extension, Cooperative Extension. All rights reserved.

Authors: Carrie Laboski is associate professor of soil science and John Peters is director of the University of Wisconsin–Madison soil testing labs in Madison and Marshfield. Both hold joint appointments with the College of Agricultural and Life Sciences, University of Wisconsin–Madison, and University of Wisconsin–Extension, Cooperative Extension. Cooperative Extension publications are subject to peer review.

University of Wisconsin–Extension, Cooperative Extension, in cooperation with the U.S. Department of Agriculture and Wisconsin counties, publishes this information to further the purpose of the May 8 and June 30, 1914, Acts of Congress. An EEO/AA employer, the University of Wisconsin–Extension, Cooperative Extension provides equal opportunities in employment and programming, including Title IX and ADA requirements. If you have a disability and require this information in an alternative format, please contact Cooperative Extension Publishing at 432 N. Lake St., Rm. 227, Madison, WI 53706; pubs@uwex.edu; or (608) 263-2770 (711 for Relay).

This publication is available from your county UW-Extension office (www.uwex.edu/ces/cty) or from Cooperative Extension Publishing. To order, call toll-free 1-877-947-7827 or visit our website at learningstore.uwex.edu.

